arXiv:0911.5235v1 [math.DS] 27 Nov 2009

HAMILTONIAN AND SMALL ACTION VARIABLES FOR PERIODIC DNLS

EVGENY L. KOROTYAEV

Abstract. We consider the defocussing NLS equation with small periodic initial condition. A new approach to study the Hamiltonian as a function of action variables is demonstrated. The problems for the NLS equation is reformulated as the problem of conformal mapping theory corresponding to quasimomentum of the Zakharov-Shabat operator. The main tool is the Löwner type equation for the quasimomentum. In particular, we determine the asymptotics of the Hamiltonian for small action variables. Moreover, we determine the gradient of Hamiltonian with respect to action variables. This gives so called frequencies and determines how the angles variables depend on the time.

Dedicated to the memory of my teacher Mikhail Sh. Birman, 1928-2009

1. Introduction

Consider the defocussing cubic non-linear Schrödinger equation (dNLS)

$$J\frac{\partial\psi}{\partial t}=-\psi_{xx}+2|\psi|^2\psi,\quad J=\begin{pmatrix}0&1\\-1&0\end{pmatrix},\qquad \psi=\begin{pmatrix}\psi_1\\ \psi_2\end{pmatrix},$$

on the circle $\mathbb{T}=\mathbb{R}/\mathbb{Z}$, i.e. $\psi(x+1,t)=\psi(x,t)$ for $x,t\in\mathbb{R}$, with the initial conditions:

$$\psi(\cdot,0)=q=\begin{pmatrix}q_1\\ q_2\end{pmatrix}\in L^2(\mathbb{T})\oplus L^2(\mathbb{T}),$$

where q_1,q_2 are real functions. The dNLS equation has the Hamiltonian H given by

$$H(q)=\frac12\int_0^1(|q'(x)|^2+|q(x)|^4)dx. \tag{1.1}$$

For functionals $E=E(q)$ and $G=G(q)$ the Poisson bracket $\{E,G\}_P$ has the form

$$\{E,G\}_P=\int_0^1\left(\frac{\partial E}{\partial q_1(x)}\frac{\partial G}{\partial q_2(x)}-\frac{\partial G}{\partial q_1(x)}\frac{\partial E}{\partial q_2(x)}\right)dx.$$

The Hamiltonian system with Hamiltonian H is given by

$$\frac{\partial\psi}{\partial t}=\{H(\psi),\psi\}_P=-J\begin{pmatrix}\frac{\partial H}{\partial\psi_1}\\ \frac{\partial H}{\partial\psi_2}\end{pmatrix}=-J(-\psi_{xx}+2|\psi|^2\psi).$$

The dNLS equation has the Hamiltonian H and other two integrals H_0 and H_1 given by

$$H_0(q)=\|q\|^2=\int_0^1(q_1(x)^2+q_2(x)^2)dx,$$

Date: November 30, 2009.
1991 *Mathematics Subject Classification.* 35Q55, (37K10, 37K20).
Key words and phrases. periodic NLS, Hamiltonian, action variables.

$$H_1(q)=\int_0^1 (q_2'(x)q_1(x)-q_1'(x)q_2(x))dx.$$

There are many papers and books devoted to the dNLS equation, see [FT], [GH]. The action-angle variables for the dNLS equation on the circle were studied by McKean-Vaninsky [MV1], [MV2], and Grebert-Kappeler-Pöschel [GKP], see also [V]. Note that the action-angle variables are studied in [VN], [Ku], [KP]. The dNLS equation admits globally defined real analytic action-angle variables $A_n, \phi_n, n\in\mathbb{Z}$ (see [GKP], [MV1], [MV2], [VN]). Recall the important identities from [K2]:

$$H_0=\|q\|^2=\sum_{n\in\mathbb{Z}} A_n, \tag{1.2}$$

$$H_1=\sum_{n\in\mathbb{Z}}(2\pi n)A_n, \tag{1.3}$$

$$H=\sum_{n\in\mathbb{Z}}(2\pi n)^2A_n+2H_0^2-U, \tag{1.4}$$

where U is some nonlinear functional given by (4.2). In fact we rewrite H_0, H_1 and the main part of H in terms of simple functions of the actions $A=(A_n)_{n\in\mathbb{Z}}$. Moreover, the following estimates from [K2] hold true:

$$0\leqslant U(A)\leqslant \frac{4}{3}\|A\|_1^2, \qquad if \quad |q|\in L^2(\mathbb{T}), \tag{1.5}$$

where $\|A\|_1=\sum_{n\in\mathbb{Z}}A_n$. Introduce the real spaces

$$\ell^p=\{f=(f_n)_{n\in\mathbb{Z}},\ \ \|f\|_p<\infty\}, \qquad \|f\|_p^p=\sum_n f_n^p<\infty,\ p\geqslant 1.$$

Note that (1.2), (1.4), (1.5), yield:
(i) $|q|\in L^2(\mathbb{T})$ iff $(A_n)_{n\in\mathbb{Z}}\in\ell^1$,
(ii) $|q'|\in L^2(\mathbb{T})$ iff $(nA_n)_{n\in\mathbb{Z}}\in\ell^1$.

In Theorem 1.1 we will show that $U(A)$ is a well defined function of $A=(A_n)_{n\in\mathbb{Z}}\in\ell^2$. We mention the Kuksin conjecture [Ku1]: *there is an estimate of the Hamiltonian H in terms of $\mathcal{P}(\|A\|_2)$ for some polynomial $\mathcal{P}(z)$.* As far as we know, no an estimate of H in terms of $\mathcal{F}(\|A\|_2)$ for some function $\mathcal{F}(z), z\geqslant 0$ has been published.

We formulate our result about the estimates.

Theorem 1.1. *The following estimates hold true:*

$$\frac{\pi}{6}\|A\|_2^2\leqslant U(A)\leqslant\frac{2\pi}{3}\sqrt{C_1}\|A\|_2^2, \qquad if \qquad (|\hat q_n|)_{n\in\mathbb{Z}}\in\ell^4, \tag{1.6}$$

where $\hat q_n=\int_0^1 q(x)e^{-i2\pi nx}dx$ *and* $C_1=\max\{2,\cosh\frac{\pi}{2}\|A\|_\infty\}$.

Remark. 1) $A\in\ell^2$ iff $(|\hat q_n|)_{n\in\mathbb{Z}}\in\ell^4$, see Sect. 2.
2) If $q\in L^{\frac{4}{3}}(\mathbb{T})$, then $(|\hat q_n|)_{n\in\mathbb{Z}}\in\ell^4$.
2) (1.6) gives $U(A)=0$ for some $A\in\ell^2$ iff $A=0$ (or $q=0$).

The Hamiltonian H depends only on the actions A. Introduce the frequencies Ω_n by

$$\Omega_n=\partial_n H,\quad n\in\mathbb{Z},\qquad \partial_n=\frac{\partial}{\partial A_n},\qquad \partial=(\partial_n)_{n\in\mathbb{Z}}. \tag{1.7}$$

The parameters Ω_n are very important, since the angle variables $\phi_n(t)$ as functions of time $t \geqslant 0$ have the form

$$\phi_n(t) = \phi_n(0) + \Omega_n t, \quad t \geqslant 0, \quad n \in \mathbb{Z}.$$

Due to (4.1) we deduce that the gradient is given by

$$\Omega_n = \partial_n H = (2\pi n)^2 + 4H_0 - \partial_n U. \tag{1.8}$$

Thus in order to study Ω_n we need to study $\partial_n U$ only, which is defined on $q \in L^2(\mathbb{T})$. Our goal is to give a new method to study Hamiltonian as a function of action variables. In this paper we reformulate the problems for the dNLS equation as the problems of the conformal mapping theory. The main technical tool is Theorem 3.1 from [KK3] about the Löwner type equation. We formulate our main result.

Theorem 1.2. *Let* $\mathscr{A} = \{A \in \ell^1 : \sum_{n\in\mathbb{Z}} A_n \leqslant \frac{1}{8^2}\}$. *Then the function* $U : \mathscr{A} \to [0,\infty)$ *has the derivative* $\partial_n U(A)$ *for each* $n \in \mathbb{Z}$, *which is continuous on* $\mathscr{A}$ *and satisfies*

$$|U(A) - \|A\|_2^2| \leqslant 4\pi\sqrt{3}\|A\|_2^3, \tag{1.9}$$

$$\|\partial U(A) - 2A\|_2 \leqslant 11\pi^2 \|A\|_\infty \|A\|_2. \tag{1.10}$$

Remark. 1) Note that the Hamiltonian $H(q)$ is defined on the functions $|q'| \in L^2(\mathbb{T})$, but we study the frequencies for the potentials $|q| \in L^2(\mathbb{T})$.
2) In order to prove (1.9) with the reminder of the third order $\|A\|_2^3$ we use the properties of the quasimomentum from [KK2], [K3],[K4] and the identity (4.1) only. Moreover, similar arguments give the Taylor series of the Hamiltonian H in terms of actions A_n for any order. We consider the simplest case to write the short paper and to formulate our approach. The main problem is to show (1.10).
3) We estimate U in terms of the norm $\|A\|_2$, which yields estimates in terms of $\|A\|_1$, since $\|A\|_2 \leqslant \|A\|_1$. In order to study the perturbation of the dNLS the estimates the Hamiltonian $H(q)$ in terms of $\|A\|_2$ are important. This is the motivation of the estimates (1.9), (1.10).
4) In the case $|q'| \in L^2(\mathbb{T})$ the Hamiltonian H is a real analytic function of $(nA_n)_{n\in\mathbb{Z}}$ and the Marchenko-Ostrovski parameters, similar to the case H_0, see Theorem 3.1.
4) In [K5] we consider the frequencies Ω_n for general case and determine their asymptotics as $n \to \pm\infty$.

We now describe the plan of the paper. In Section 2 we recall the needed results from [KK2], [KK3], [K4], [K3] and [MO] about the quasimomentum $k(z,h)$ as the function of two variables z, h: z is a spectral variable and $h \in \ell^2$ is the Marchenko-Ostrovski parameter. Note that for each $h \in \ell^2$ the function $k(\cdot,h) : \mathcal{K}(h) \to \mathcal{Z}$ is the conformal mapping (see definitions of $\mathcal{K}(h), \mathcal{Z}$ and $k(\cdot,h)$ in (2.4)). In Theorem 3.1 from [KK3] we recall the needed properties of the function $z(k,h)$ (for fix $h \in \ell^2$ the function $z(k,h)$ is an inverse for $k(z,h)$) as the function of quasimomentum k and $h \in \ell^2$ and formulate the Löwner type equation (3.4) for the conformal mapping $z(\cdot,h) : \mathcal{K}(h) \to \mathcal{Z}$. In Lemma 3.2 from [KK3] we describe the properties of the actions $A_n(h)$ as functions of the Marchenko-Ostrovski parameters $h \in \ell^2$ and present the exact formulas for $\frac{\partial}{\partial h_m} A_n(h), n, m \in \mathbb{Z}$. In the proof of Theorem 1.1 we use the identity $\omega_m = \sum_{n\in\mathbb{Z}} X_{m,n}\widetilde{\omega}_n$, where $\widetilde{\omega}_m = \frac{\partial U}{\partial h_m^2}$ and $X_{m,n} = \frac{\partial h_m^2}{\partial A_n}, m \in \mathbb{Z}$. In Lemma 3.3 we show that operator $X : \ell^2 \to \ell^2$ with the matrix $X_{m,n}$ satisfies $X = I + o(1)$ as $A \to 0$. In Lemma 4.1 we determine $\widetilde{\omega}_n = 2A_n + o(\|A\|)$ as $A \to 0$. Then roughly speaking we determine asymptotics $\omega_n = \frac{\partial U}{\partial A_n}$.

2. **Preliminaries**

The dNLS equation is integrable and admits a Lax-pair formalism, see [ZS], [FT]. Consider the corresponding Zakharov-Shabat operator T_{zs} acting in $L^2(\mathbb{R})\oplus L^2(\mathbb{R})$ and given by

$$T_{zs}=J\frac{d}{dx}+\mathcal{Q},\qquad \mathcal{Q}=\begin{pmatrix} q_1 & q_2\\ q_2 & -q_1\end{pmatrix}, \tag{2.1}$$

where $|q|\in L^1(0,1)$. This operator is essentially self-adjoint on the domain $\mathscr{D}=\{f,f',f''\in L^2(\mathbb{R})\oplus L^2(\mathbb{R})\}$, (see [LM]). The spectrum of T_{zs} is purely absolutely continuous and is union of spectral bands $\sigma_n, n\in\mathbb{Z}$, where

$$\sigma_n=[z_{n-1}^+,z_n^-],\ \dots<z_{2n-1}^-\leqslant z_{2n-1}^+<z_{2n}^-\leqslant z_{2n}^+<\dots,\quad and\quad z_n^\pm=n\pi+o(1)\ \text{as}\,|n|\to\infty.$$

The intervals σ_n and σ_{n+1} are separated by gap $g_n=(z_n^-,z_n^+)$ with the length $|g_n|\geqslant 0$. If a gap g_n is degenerate, i.e., $|g_n|=0$, then the corresponding segments σ_n,σ_{n+1} merge. We use the Zakharov-Shabat equation for a vector -function f:

$$Jf'+\mathcal{Q}f=zf,\quad z\in\mathbb{C},\qquad f=\begin{pmatrix} f_1\\ f_2\end{pmatrix}, \tag{2.2}$$

where f_1,f_2 are functions of $x\in\mathbb{R}$. Here and below $(\,')=\partial/\partial x$. The boundary value problem with Eq. (2.2) and with the condition $f(0)=f(1)$ is called periodic and the boundary value problem (2.2) with the condition $f(0)=-f(1)$ is called antiperiodic. Here $z_{2n}^\pm, n\in\mathbb{Z}$ are the eigenvalues of the periodic problem and $z_{2n+1}^\pm, n\in\mathbb{Z}$ are the eigenvalues of the anti-periodic problem. Define the 2×2-matrix valued fundamental solution $\Psi=\Psi(x,z)$ by

$$J\frac{d}{dx}\Psi+\mathcal{Q}\Psi=z\Psi,\quad \Psi(0,z)=\begin{pmatrix} 1 & 0\\ 0 & 1\end{pmatrix},\quad z\in\mathbb{C}. \tag{2.3}$$

Introduce the Lyapunov function $\Delta(z)$ by

$$\Delta(z)=\frac{1}{2}\mathrm{Tr}\Psi(1,z),\quad z\in\mathbb{C}.$$

The function Δ is entire and $\Delta(z_n^\pm)=(-1)^n$ for all $n\in\mathbb{Z}$ (see e.g. [LS]).

We recall results which is crucial for the present paper. For each $q\in L^1(\mathbb{T})$ there exists a unique conformal mapping (the quasimomentum) $k:\mathcal{Z}\to\mathcal{K}(h)$ with asymptotics $k(z)=z+o(1)$ as $|z|\to\infty$ (see Fig. 1 and 2) and such that (see [MO] and [Mi1], [Mi2], [KK2])

$$\begin{aligned}&\cos k(z)=\Delta(z),\quad z\in\mathcal{Z}=\mathbb{C}\setminus\cup\overline{g}_n,\quad and\\ &\mathcal{K}(h)=\mathbb{C}\setminus\cup\Gamma_n,\quad \Gamma_n=(\pi n-i|h_n|,\pi n+i|h_n|),\qquad h_n\in\mathbb{R},\\ &|h_n|\geqslant 0\qquad \text{is defined by the equation}\qquad \cosh|h_n|=(-1)^n\Delta(z_n)\geqslant 1.\end{aligned} \tag{2.4}$$

Here Γ_n is the vertical cut and recall that $z_n\in[z_n^-,z_n^+]$ and $\Delta'(z_n)=0$. Moreover, we have $(|h_n|)_{n\in\mathbb{Z}}\in\ell^2$ iff $q\in L^2(0,1)$ (and $(n|h_n|)_{n\in\mathbb{Z}}\in\ell^2$ iff $q'\in L^2(\mathbb{T})$), see [K1], [K2].

Recall the properties of the conformal mapping $k=u(z)+iv(z), z\in\mathcal{Z}$ from [MO] or [KK2]:

Lemma 2.1. *Let $h\in\ell^\infty$. Then the quasimomentum $k=u(z)+iv(z), z\in\mathcal{Z}$ satisfies:*
1) $v(z)\geqslant\operatorname{Im}z>0$ and $v(z)=-v(\overline{z})$ for all $z\in\mathbb{C}_+=\{\operatorname{Im}z>0\}$.
2) $v(z)=0$ for all $z\in\sigma_n, n\in\mathbb{Z}$.

3) If some $g_n \neq \emptyset, n \in \mathbb{Z}$, then the function $v(z+i0) > 0$ for all $z \in g_n$, and $v(z+i0)$ has a maximum at $z_n \in g_n$ such that $v(z_n+i0) = |h_n|, v'(z_n) = 0$, see Fig. 3, and $\Delta'(z_n) = 0$ and

$$v(z+i0) = -v(z-i0) > 0, \qquad \frac{v'(z+i0)}{z_n - z} > 0, \qquad v''(z+i0) < 0, \qquad \text{all } z \in g_n \neq \emptyset, \tag{2.5}$$

$$|g_n| \leqslant 2|h_n|. \tag{2.6}$$

4) $u'(z) > 0$ on all (z_{n-1}^+, z_n^-) and $u(z) = \pi n$ for all $z \in g_n \neq \emptyset, n \in \mathbb{Z}$.
5) The function $k(z)$ maps a horizontal cut (a "gap") $[z_n^-, z_n^+]$ onto vertical cut Γ_n and a spectral band σ_n onto the segment $[\pi(n-1), \pi n]$ for all $n \in \mathbb{Z}$.

We emphasize that the introduction of the quasi-momentum $k(\cdot)$ provides a natural labeling of all gaps g_n (including the empty ones!) by demanding that $k(\cdot)$ maps the cut (a "gap") $[z_0^-, z_0^+]$ on the vertical cut $\Gamma_0 = (-ih_0, ih_0)$. This determination of a fixed reference point will be of important later on. Let $z(\cdot) = k^{-1} : \mathcal{K}(h) \to \mathcal{Z}$ be the inverse mapping for $k : \mathcal{Z} \to \mathcal{K}(h)$. Below we will sometimes write $z(k,h), , ..,$ instead of $z(k), , ..,$ when several h are being dealt with. Note, that for fixed $k \in \mathcal{K}(h)$ the function $z(k,h)$, $h \in \ell^2_{\mathbb{R}}$ is **even with respect to each variable** $h_n \in \mathbb{R}, n \in \mathbb{Z}$. We use $h_n \in \mathbb{R}$, since it is more convenient for us than $h_n \in [0,\infty)$.

For any nondegenerate gap g_n the function $k(\cdot,h)$ has an analytic continuation (from above or from below) across the interval g_n. This suffices to extend the function $-i(k(\cdot,h) - \pi n)$ by the symmetry. Similarly the function $z(\cdot,h)$ has an analytic continuation (from left or from right) across the vertical cut $(\pi n - i|h_n|, \pi n + i|h_n|)$ by the symmetry.

In spirit, such result goes back to the classical Hilbert Theorem (for a finite number of cuts, see e.g. [J]) in the conformal mapping theory. A similar theorem for the Hill operator is technically more complicated (there is a infinite number of cuts) and was proved by Marchenko-Ostrovski [MO]. The proof of Misura for the Zakharov-Shabat system follows the general idea from [MO]. For additional properties of the conformal mapping we also refer to our previous papers [KK2], [KK4], [K2].

A lot of papers are devoted to the inverse problems for the operator T_{zs}. Misura [Mi1], [Mi2] extended the results of [MO] to the periodic Zakharov-Shabat operator (in terms of heights $h = (h_n)_{n\in\mathbb{Z}} \in \ell^2$ of vertical cuts $\Gamma_n, n \in \mathbb{Z}$, see Sect.2). The author [K1] reproved the results of Misura [Mi1], [Mi2] by the direct method [KK1]. The gap length mappings were considered in [BGGK], [GG], [K2]. A priori estimates (two sided) of H_0, H in terms of actions, gap lengths, Marchenko-Ostrovski parameters, etc. were obtained in [KK2], [KK3], [K1]-[K4]. There are papers devoted to the integrals H, H_0, H_1, see [KK2], [KK3], [K1]-[K4], where various both identities and estimates of integrals in terms of gap lengths of the Zakharov-Shabat operator, actions variables, the Marchenko-Ostrovski parameters (so-called heights h_n, see Sect. 2) were obtained.

Note that if $q \in L^2(\mathbb{T})$, then $v \in L^1(\mathbb{T}) \cap L^\infty(\mathbb{R})$, see (2.8).

The quasimomentum $k(\cdot)$ has has asymptotics

$$k(z) = z - \frac{Q_0 + o(1)}{z} \quad \textit{if} \;\; q \in L^2(\mathbb{T}) \quad \textit{and} \quad k(z) = z - \frac{Q_0}{z} - \frac{Q_1}{z^2} - \frac{Q_2 + o(1)}{z^3} \quad \textit{if} \;\; q' \in L^2(\mathbb{T})$$

as $z \to i\infty$, see [K2], where the functionals $Q_j(h), (n^{\frac{j}{2}} h_n)_{n\in\mathbb{Z}} \in \ell^2, j = 0,1,2$ are given by

$$Q_j(h) = \frac{1}{\pi} \int_{\mathbb{R}} z^j v(z+i0,h)dz \geqslant 0, \qquad j = 0,1,2, \quad k(z) = u(z) + iv(z). \tag{2.7}$$



FIGURE 1. The domain $\mathscr{Z}=\mathbb{C}\setminus\cup\bar{g}_n$, where $g_n=(z_n^-,z_n^+)$



FIGURE 2. The domain $K(h)=\mathbb{C}\setminus\cup\Gamma_n$, where $\Gamma_n=(\pi n-ih_n,\pi n+ih_n)$



FIGURE 3. The graph of $v(z+i0)$, $z\in g_n\cup\sigma_n\cup\sigma_{n+1}$ and $|h_n|=v(z_n+i0)>0$

Recall the following identities from [KK1]

$$H_0=2Q_0(h)=\frac{1}{\pi}\iint_{\mathbb{C}}|z'(k,h)-1|^2dudv,\qquad H_1=4Q_1(h),\qquad H_2=8Q_2(h),\quad(2.8)$$

where $k=u+iv$. The functions $Q_j(h)$ are even with respect to each variable h_n, and then due to (3.5), $Q_j(h)$ are the functions of $A=(A_n)_{n\in\mathbb{Z}}$.

Recall estimates for the periodic Zakharov-Shabat operator (see Corollary 2.3, [K3])

$$\frac{1}{2}\|q\|\leqslant\|h\|_2\leqslant 3(1+\|q\|)^{\frac{1}{2}}\|q\|,\qquad(2.9)$$

$$\frac{1}{2}\|\widetilde{g}\|_2 \leqslant \|q\| \leqslant 2\|\widetilde{g}\|_2(1+\|\widetilde{g}\|_2), \qquad \widetilde{g} = (|g_n|)_{n\in\mathbb{Z}}, \tag{2.10}$$

$$\|\eta\|_2 \leqslant 16\min\Big\{\|q\|,\ \|h\|_2,\ \|\widetilde{g}\|_2(1+\|\widetilde{g}\|_2)\Big\}, \tag{2.11}$$

where $\eta = (\eta_n)_{n\in\mathbb{Z}}$ and $\eta_n = \pi - |\sigma_n| \geqslant 0$, see [K3].

The dNLS equation admits globally defined real analytic action-angle variables $A_n, \phi_n, n \in \mathbb{Z}$ (see [GKP], [MV1], [MV2], [VN]), where the action variables A_n are given by

$$A_n = -\frac{1}{\pi i}\int_{\chi_n} z\frac{\Delta'(z)}{\sqrt{1-\Delta^2(z)}}dz \geqslant 0, \tag{2.12}$$

see [FM], where $\chi_n \subset \mathbb{C}$ is a counterclockwise circuit around the gap g_n only and the branch of $\sqrt{1-\Delta^2(z)}$ is defined by $\sqrt{1-\Delta^2(z+i0)} > 0$ for all $z \in (z_0^+, z_1^-)$.

Using the identity $\Delta(z) = \cos k(z)$ and integration by parts we rewrite the action A_n in terms of the quasimomentum (see [KK3])

$$A_n(h) = -\frac{1}{\pi i}\int_{c_n} k(z,h)dz = \frac{2}{\pi}\int_{g_n} v(z,h)dz \geqslant 0, \quad v = \operatorname{Im} k(z), \qquad n \in \mathbb{Z}. \tag{2.13}$$

Here and below the function $v(z,h)$ in the integral $\int_{g_n} v(z,h)dz$ on the gap $g_n \neq \emptyset$ is given by $v(z,h) = v(z+i0,h) > 0, z \in g_n$. Below we need the following estimates from [KK1]:

$$\max\left\{\frac{|g_n|^2}{4}, \frac{|g_n||h_n|}{\pi}\right\} \leqslant A_n = \frac{2}{\pi}\int_{g_n} v(z)dz \leqslant \frac{|g_n||h_n|}{\pi} \leqslant \frac{2|h_n|^2}{\pi}, \quad \text{all } n \in \mathbb{Z}. \tag{2.14}$$

We need the important identity from [KK2] (see Fig. 3)

$$v(x) = v_n(x)(1+Y_n(x)), \quad Y_n(x) = \frac{1}{\pi}\int_{\mathbb{R}\setminus g_n} \frac{v(t)dt}{|t-x|\,v_n(t)} \geqslant 0, \quad \text{all} \quad x \in g_n = (z_n^-, z_n^+),$$
$$v_n(x) = |(x-z_n^+)(x-z_n^-)|^{\frac{1}{2}} = |r_n^2 - (z-z_n^0)^2|^{\frac{1}{2}}, \qquad z_n^0 = \frac{z_n^- + z_n^+}{2}, \quad r_n = \frac{|g_n|}{2}. \tag{2.15}$$

For each $n \in \mathbb{Z}$ there exists the unique points $z_n \in g_n$ such that

$$v(z_n + i0) = |h_n| = v_n(z_n)(1+Y_n(z_n)). \tag{2.16}$$

Define $s = \min_{n\in\mathbb{Z}}|\sigma_n|$ and the sequences $M_n, ..,$ and the sequence $S = (S_m)_{m\in\mathbb{Z}}$ by

$$M_n = \max_{z\in g_n} Y_n(z), \quad \dot{M}_n = \max_{z\in g_n}|Y_n'(z)|, \quad \ddot{M}_n = \max_{z\in g_n}|Y_n''(z)|, \quad S_m = \frac{1}{2}\sum_{n\neq m}\frac{A_n}{s^2(n-m)^2}.$$

Lemma 2.2. *Let $\|q\| \leqslant \frac{1}{8}$. Then for each $n \in \mathbb{Z}$ the following estimates hold true:*

$$s = \min|\sigma_n| \geqslant 1, \quad \sup|g_n| \leqslant \frac{1}{4}, \tag{2.17}$$

$$\sup_{z\in g_m}\frac{1}{\pi}\int_{g\setminus g_m}\frac{v(t,h)dt}{(t-z_m)^2} \leqslant S_m, \tag{2.18}$$

$$M_n \leqslant S_n, \qquad \dot{M}_n \leqslant S_n, \qquad \ddot{M}_n \leqslant S_n, \tag{2.19}$$

$$\|S\|_\infty \leqslant \frac{H_0}{2} \leqslant \frac{1}{128}, \qquad \|S\|_2 \leqslant \frac{\pi^2}{6}\|A\|_2, \qquad \|S\|_1 \leqslant \frac{\pi^2}{6}H_0 \leqslant \frac{\pi^2}{6\cdot 64}. \tag{2.20}$$

Proof. The estimates (2.11) gives $0 \leqslant \pi - |\sigma_n| \leqslant 2$ and then $s \geqslant 1$.

The estimates (2.10) gives $\sum |g_n|^2 \leqslant 4\|q\|^2 \leqslant \frac{1}{16}$, which yields (2.17).

Using $s \geqslant 1$ and (1.2) we get $S_m \leqslant \frac{1}{2}\sum_{n\neq m} A_n = \frac{H_0}{2}$. The definition of S_m gives

$$\sum_m S_m^2 \leqslant \frac{1}{4}\sum_m \sum_{p\neq m} \frac{1}{(p-m)^2} \sum_{n\neq m} \frac{A_n^2}{(n-m)^2} = \frac{\pi^4}{6^2}\|A\|_2^2,$$

since $\sum_{n>0} \frac{1}{n^2} = \frac{\pi^2}{6}$. Similar arguments yield the last estimate in (2.19) and (2.18), (2.20) . ■

Lemma 2.3. *Let $h \in \ell^\infty$. Then for each $n \in \mathbb{Z}$ the following estimates hold true:*

$$2|h_n| \leqslant |g_n|(1 + M_n), \tag{2.21}$$

$$0 \leqslant A_n - \frac{|g_n|^2}{4} \leqslant \frac{|g_n|^2}{4} M_n, \tag{2.22}$$

$$|h_n - \nu_n| \leqslant 4h_n M_n + h_n \frac{|g_n|^2}{4}\biggl(3(1 + \frac{|g_n|}{2})\dot{M}_n^2 + \ddot{M}_n\biggr), \tag{2.23}$$

$$|2h_n - |g_n|(1 + Y_n(z_n)| \leqslant \frac{|g_n|^3}{8}\dot{M}_n^2, \tag{2.24}$$

$$|A_n - \frac{|g_n| h_n}{4}| \leqslant \frac{|g_n|^4}{2^7}(\ddot{M}_n + 6\dot{M}_n^2), \tag{2.25}$$

$$|z_n - z_n^0| \leqslant \frac{|g_n|^2}{4}\dot{M}_n, \qquad \textit{where} \qquad z_n^0 = \frac{z_n^- + z_n^+}{2}. \tag{2.26}$$

Proof. Estimates (2.24)-(2.26) were proved in [K4] (see Theorem 1.3 and 1.4 in [K4]).

Estimate (2.16) implies (2.21). Using (2.15) we obtain

$$A_n = \frac{2}{\pi}\int_{g_n} v_n(x)(1 + Y_n(x))dx = \frac{|g_n|^2}{4} + \frac{2}{\pi}\int_{g_n} v_n(x) Y_n(x) dx,$$

which gives (2.22). We will show (2.23). Let $r = \frac{|g_n|}{2}$ and $\alpha_n = \frac{\nu_n}{h_n}$. We need the estimate

$$\left|\frac{r^2}{h_n^2} - \alpha_n\right| \leqslant r^2 C_0, \quad C_0 = (3 + r)\dot{M}_n^2 + \ddot{M}_n, \tag{2.27}$$

(see Lemma 2.1 in [K4]). Then we obtain

$$|\alpha_n - 1| \leqslant |\alpha_n - (r^2/h_n^2)| + |(r^2/h_n^2) - 1| \leqslant r^2 C_0 + 2|(r/h_n) - 1|. \tag{2.28}$$

The estimate (2.24) implies

$$\left|\frac{r}{h_n} - 1\right| \leqslant \left|\frac{r}{h_n} - \frac{1}{1 + Y_n(z_n^0)}\right| + \frac{Y_n(z_n^0)}{1 + Y_n(z_n^0)} \leqslant r^3 \dot{M}_n^2 + 2M_n. \tag{2.29}$$

Then $|\alpha_n - 1| \leqslant 4M_n + r^2(3(1 + r)\dot{M}_n^2 + \ddot{M}_n)$, which yields (2.23). ■

3. The action $A_n(h)$ as a function of h

Define the ball $\mathcal{B}^p(r) = \{\eta : \|\eta\|_p \leqslant r\} \subset \ell^p, r > 0$. Let ℓ^p_C be the complexification of the space ℓ^p. In the complex space ℓ^p_C the corresponding ball is denoted by $\mathcal{B}^p_C(r) \subset \ell^p_C$. Define the strip

$$\mathfrak{J}_\beta = \{\eta \in \ell^2_C : \|\operatorname{Im}\eta\|_2 < \beta\}, \qquad \beta > 0. \tag{3.1}$$

Recall that for fixed $k \in \mathbb{C}$ the function $z(k,h),\ h \in \ell^2_{\mathbb{R}}$ is even with respect to each variable $h_n \in \mathbb{R}$. Below we need some results about the quasimomentum (see Theorem 2.3, 2.5 from [KK3]).

Theorem 3.1. *i) The functional $Q_0 : \ell^2 \to \mathbb{R}_+$ has an analytic continuation into some strip $\mathfrak{J}_\beta, \beta > 0$ and the gradient is given by:*

$$\nabla_n Q_0(h) = \nu_n(h) = (-1)^{n-1}\frac{\sinh h_n}{\Delta''(z_n)}, \qquad \text{all} \quad n \in \mathbb{Z}, \quad \nabla_n = \frac{\partial}{\partial h_n}. \tag{3.2}$$

ii) There exist $\varepsilon > \beta > 0$ such that for any fixed real $h \in \ell^2$ the function $z(k, h+\eta)$ has the analytic extension from $(k,\eta) \in \mathcal{K}(h,\varepsilon) \times \mathcal{B}^2(\beta)$ into the domain $K(h,\varepsilon) \times \mathcal{B}^2_C(\beta)$, where

$$\mathcal{K}(h,\varepsilon) = \{k \in \mathbb{C} : \ \operatorname{dist}(k, \cup\Gamma_n) > \varepsilon\} \subset \mathcal{K}(h).$$

Moreover, for each $n \in \mathbb{Z}$ the derivatives are given by (the Löwner type equation)

$$\nabla_n z(k,h) = 0, \quad h_n = 0, \qquad k \neq \pi n, \ k \in \mathcal{K}(h), \tag{3.3}$$

$$\nabla_n z(k,h) = \frac{\nu_n}{z(k,h) - z_n(h)}, \quad h_n \neq 0, \qquad k \neq \pi n \pm i h_n, \ k \in \mathcal{K}(h). \tag{3.4}$$

iii) The mapping $a : \ \ell^2_j \to \ell^2_j$ for all $j \in [0,\infty)$ given by

$$h \to a = (a_n)_{n\in\mathbb{Z}}, \qquad a_n(h) = |A_n(h)|^{\frac{1}{2}} \operatorname{sign} h_n, \tag{3.5}$$

is a real analytic isomorphism of $\ell^2_j = \{f = (f_n)_{n\in\mathbb{Z}} : \sum_{n\in\mathbb{Z}}(1+n^2)^j f_n^2 < \infty\}$ onto itself.

Proof. The statement (i) was proved in [KK3] with ν_n given by

$$\nu_n = \begin{cases} |k''(z_n,h)|^{-1} \operatorname{sign} h_n, & if \quad |g_n| > 0, \\ 0, & if \quad |g_n| = 0, \end{cases}. \tag{3.6}$$

Using $\Delta(z) = \cos k$ we obtain $\Delta'(z) = -k'(z)\sin k$ and thus $\Delta''(z_n) = -k''(z)\sin k(z_n)$, since $\Delta'(z_n) = 0$. This and $\sin k(z_n) = i(-1)^n \sinh h_n =$ gives $\Delta''(z_n) = v''(z_n)(-1)^n \sinh h_n$, which yields (3.3), (3.4). All other statements were proved in [KK3]. ■

If $h \in \ell^\infty_{\mathbb{R}}$, then for each $n \in \mathbb{Z}$ the following estimates from [KK3] hold true:

$$\nu_n^2 \leqslant h_n^2 \leqslant H_0, \qquad where \qquad \nu_n(h) = (-1)^{n-1}\frac{\sinh h_n}{\Delta''(z_n)}, \tag{3.7}$$

and recall that $z_n \in [z_n^-, z_n^+]$ is the zero of the function $\Delta'(z)$.

Note that (3.4) is the Löwner type equation for the quasimomentum.

Introduce the contours c_n around Γ_n and χ_n around g_n by

$$c_n = \{k : \operatorname{dist} \ (k,\Gamma_n) = \pi/4\} \subset \mathcal{K}(h), \qquad \chi_n = z(c_n,h) \subset \mathcal{Z}. \tag{3.8}$$

Below we need some results about the action variable A_n (see Lemma 7.6 from [KK3]).

Lemma 3.2. *Each action* $A_n : \ell^2 \to [0,\infty), n \in \mathbb{Z}$ *has an analytic continuation into some strip* $\mathfrak{J}_\beta, \beta > 0$ *by*

$$A_n(h) = -\frac{1}{\pi i}\int_{c_n} z(k,h)dk. \tag{3.9}$$

Moreover, the derivatives $\frac{\partial A_n}{\partial h_m}, m \in \mathbb{Z}$ *have the following forms:*

$$\nabla_m A_n(h) = \frac{\nu_m}{\pi i}\int_{c_n} \frac{dk}{z(k,h) - z_m}, \qquad h \in \ell^2, \quad \nabla_m = \frac{\partial}{\partial h_m}, \tag{3.10}$$

where $\nu_m = \nu_m(h), z_m = z_m(h)$, *and if* $h \in \ell^2$, *then*

$$\nabla_n A_n(h) = 2\nu_n + \frac{2\nu_n}{\pi}\int_{g\setminus g_n} \frac{v(z,h)dz}{(z-z_n)^2}, \qquad g = \cup g_s, \tag{3.11}$$

$$\nabla_m A_n(h) = \frac{2\nu_m}{\pi}\int_{g_n} \frac{v'(z,h)dz}{z_m - z}, \tag{3.12}$$

$$\nabla_m A_n(h) = -\frac{2\nu_m}{\pi}\int_{g_n} \frac{v(z,h)dz}{(z-z_m)^2}, \quad m \neq n. \tag{3.13}$$

Remark. Recall that that $\frac{v'(z+i0)}{z_m - z} > 0$ for all $z \in g_m$ and the identities (3.11), (3.12) and (2.18) give

$$\frac{1}{\pi}\int_{g_m} \frac{v'(z)dz}{z_m - z} = 1 + \frac{1}{\pi}\int_{g\setminus g_m} \frac{v(z)dz}{(z-z_m)^2} \leqslant 1 + S_m, \qquad \text{all } m \in \mathbb{Z}, \tag{3.14}$$

Define the operator $F : \ell^2 \to \ell^2$ by

$$(Ff)_m = \sum_{n\in\mathbb{Z}} F_{m,n} f_n, \qquad F_{m,n} = \frac{\partial A_n(h)}{\partial h_m^2}, \quad f = (f_n)_{n\in\mathbb{Z}}. \tag{3.15}$$

Introduce the sequence $\alpha_n = \frac{\nu_n}{2h_n}, n \in \mathbb{Z}$. Due to Lemma 3.2 we obtain

$$F_{m,n} = -\frac{\alpha_m}{\pi}\int_{g_n} \frac{v(z,h)dz}{(z-z_m)^2}, \qquad \alpha_m = \frac{\nu_m}{h_m}, \qquad m \neq n, \tag{3.16}$$

$$F_{n,n} = \alpha_n + \frac{\alpha_n}{2\pi}\int_{g\setminus g_n} \frac{v(z,h)dz}{(z-z_n)^2}, \qquad g = \cup g_s. \tag{3.17}$$

Lemma 3.3. *Let* $\|q\| \leqslant \frac{1}{8}$. *Then* $F - I_{id}$ *is the Hilbert-Schmidt operator and satisfies*

$$|F_{m,n}| \leqslant \frac{A_n}{2(n-m)^2}, \qquad m \neq n, \tag{3.18}$$

$$|F_{n,n} - 1| \leqslant 5S_n, \tag{3.19}$$

$$\|F - I_{id}\|_{HS} \leqslant \pi\|q\|^2 \leqslant \frac{\pi}{64}, \tag{3.20}$$

$$|\alpha_n - 1| \leqslant 5S_n, \tag{3.21}$$

where $\|\cdot\|_{HS}$ *is a Hilbert-Schmidt norm and the operator* F *has an inverse.*

Proof. If $m \neq n$, then using (2.17), (3.7) we obtain

$$|F_{m,n}| \leqslant \frac{1}{\pi}\int_{g_n} \frac{v(x,h)dx}{(n-m)^2} = \frac{A_n}{2(n-m)^2}.$$

We show (3.19). Using (2.23) and Lemma 2.2 we obtain

$$|\alpha_n - 1| \leqslant S_n\biggl(4 + \frac{1}{64}\Bigl((3+\frac{3}{8})S_n + 1\Bigr)\biggr) \leqslant \frac{9}{2}S_n,$$

which gives (3.21), and $\frac{\alpha_n}{2\pi}\int_{g\setminus g_n}\frac{v(t,h)dt}{(t-z_n)^2} \leqslant \frac{S_n}{2}$, which yields (3.19).

Combining (3.18), (3.19) and using (2.20), (1.2), we obtain

$$\|F - I_{id}\|_{HS}^2 \leqslant 25\|S\|_2^2 + \sum_m\sum_{n\neq m}\frac{A_n^2}{4(n-m)^4}$$

$$\leqslant 25\|S\|_1\|S\|_\infty + \frac{\pi^2}{12}\|A\|_\infty^2 \leqslant \pi^2\|q\|^4\Bigl(\frac{25}{48}+\frac{1}{12}\Bigr) \leqslant \pi^2\|q\|^4 \leqslant \frac{\pi^2}{8^4},$$

which gives (3.20). ■

4. **Proof of main Theorem**

Recall that the Hamiltonian has the following form (see [K2])

$$H = \sum_{n\in\mathbb{Z}}(2\pi n)^2 A_n + 2H_0^2 - U, \qquad U(A(a)) = V(h(a)), \tag{4.1}$$

where $V(h)$ is defined by

$$V(h) = \frac{8}{3\pi}\int_{\cup g_n} v^3(z+i0,h)dz \geqslant 0, \qquad v = \operatorname{Im} k(z), \tag{4.2}$$

and $h(a)$ is an inverse mapping for $h \to a = (a_n)_{n\in\mathbb{Z}},\ a_n(h) = |A_n(h)|^{\frac{1}{2}}\operatorname{sign} h_n$ (see (3.5)) and $A(a)$ is defined by $A_n = a_n^2, n \in \mathbb{Z}$. Recall that $a \to h$ is a real analytic isomorphism of ℓ^2 onto itself (see Theorem 3.1). In fact in order to get the properties of $U(A)$ we work with the functional $V(h)$ and below we study how our functions depend on the Marchenko-Ostrovski parameter $h \in \ell^2$.

We define the functions $V_n(h), h \in \ell^2$, by

$$V(h) = \frac{8}{3\pi}\int_{\cup g_n} v^3(z,h)dz = \sum_{n\in\mathbb{Z}} V_n, \qquad V_n(h) = \frac{8}{3\pi}\int_{g_n} v^3(z,h)dz \geqslant 0. \tag{4.3}$$

Recall that $v(z+i0) > 0$ for all $z \in g_n \neq \emptyset$ and v is defined by the equation $\cosh v(z) = (-1)^n\Delta(z) \geqslant 1, z \in g_n$. Note that for fix $k \in \mathcal{K}(h)$ the function $k(z,h)$ is even with respect to each variable $h_n, n \in \mathbb{Z}$, and then due to (4.1), $V(h(a))$ is the functions of $A = (A_n)_{n\in\mathbb{Z}}$.

Using $v^2(z) \leqslant h_n^2, z \in g_n$ (see (2.15)) we obtain the simple estimate

$$V_n(h) \leqslant \frac{8h_n^2}{3\pi}\int_{g_n} v(z+i0,h)dz = \frac{4h_n^2 A_n}{3}, \qquad \textit{all} \qquad n \in \mathbb{Z}. \tag{4.4}$$

Consider now the function $V_n : \ell^2 \to [0,\infty)$, which has good properties.

Lemma 4.1. *i) Each $V_n, n \in \mathbb{Z}$ has an analytic extension from ℓ^2 into some $\mathfrak{J}_\beta, \beta > 0$ by*

$$V_n = \frac{4i}{\pi}\int_{c_n}(k-\pi n)^2 z(k,h)dk. \tag{4.5}$$

Moreover, if $h \in \ell^2$, then their gradients are given by

$$\nabla_m V_n(h) = \frac{4i\nu_m(h)}{\pi}\int_{c_n}\frac{(k-\pi n)^2}{z(k,h)-z_m(h)}dk = \frac{8\nu_m}{\pi}\int_{g_n}\frac{v^2(z)v'(z)}{z_m-z}dz, \tag{4.6}$$

$$\nabla_m V_n(h) = -\frac{8\nu_m}{3\pi}\int_{g_n}\frac{v^3(z)dz}{(z-z_m)^2}, \quad if \quad m \neq n, \tag{4.7}$$

$$\nabla_n V_n(h) = \frac{8\nu_n}{\pi}\int_{g_n}\frac{v'(z)}{z_n-z}v^2(z)dz \geqslant 0, \tag{4.8}$$

where $\nu_m = \nu_m(h), z_m = z_m(h)$ are real analytic on ℓ^2 and $v(z) = v(z+i0,h), z \in g_n$.
ii) The estimates (1.5), (1.9) and the following estimates hold true:

$$|\nabla_n V_n(h)| \leqslant 9|h_n|^3, \qquad and \qquad |\nabla_m V_n(h)| \leqslant \frac{4}{3}\frac{|h_m|h_n^2 A_n}{(n-m)^2}, \quad if \quad m \neq n. \tag{4.9}$$

Proof. i) Using $v(z+i0) = -v(z-i0)$ for all $z \in g_n$ and the integration by parts, we obtain

$$V_n = -\frac{4i}{3\pi}\int_{\chi_n}(k-\pi n)^3 dz = \frac{4i}{\pi}\int_{c_n}(k-\pi n)^2 z(k)dk.$$

The last identity and (3.4) give

$$\nabla_m V_n(h) = \frac{4i\nu_m}{\pi}\int_{c_n}\frac{(k-\pi n)^2}{z(k)-z_m}dk = \frac{4i\nu_m}{\pi}\int_{\chi_n}\frac{(k-\pi n)^2}{z-z_m}k'(z)dz,$$

which yields (4.6), (4.8). Integration by parts implies (4.7). Note that $z_m = z_m(h)$ is real analytic on ℓ^2, see [K1].

ii) We will show (1.9). Using (2.15) we obtain

$$V_n = \frac{8}{3\pi}\int_{g_n}v^3(z)dz = \frac{8}{3\pi}\int_{g_n}v_n^3(z)(1+Y_n(z))^3 dz = I_1 + I_2, \tag{4.10}$$

$$v_n(z) = |r^2-(z-z_n^0)^2|^{\frac{1}{2}}, \qquad z_n^0 = \frac{z_n^- + z_n^+}{2}, \qquad r = \frac{|g_n|}{2}.$$

Now we calculate the first integral:

$$I_1 = \frac{8}{3\pi}\int_{g_n}v_n^3(z+i0)dz = \frac{8}{3\pi}\int_{-r}^{r}|r^2-t^2|^{\frac{3}{2}}dt = r^4. \tag{4.11}$$

Using (4.10),(4.11) and $I_2 \geqslant 0$ we obtain

$$V \geqslant \frac{1}{18}\sum |g_n|^4. \tag{4.12}$$

Using (2.22) we deduce that

$$0 \leqslant A_n^2 - r^4 = (A_n - r^2)(A_n + r^2) \leqslant r^2 M_n 2A_n \leqslant 2A_n^2 M_n. \tag{4.13}$$

We estimate the second integral by

$$0 \leqslant I_2 = V_n - r^4 = \frac{8}{3\pi}\int_{g_n}v_n^3(z+i0)((1+Y_n(z))^3-1)dz$$

$$\leqslant M_n(1+M_n)^2\frac{8}{\pi}\int_{g_n} v_n^3(z+i0)dz = 3r^4M_n(1+M_n)^2 \leqslant 12r^4M_n,$$

since $Y_n(z) > 0$ for all $z \in g_n \neq 0$ and $M_n \leqslant 1$, $r^4 \leqslant A_n^2$. Thus combine the last estimates, we get (1.9), since (2.19) and $\sum_{n>0}\frac{1}{n^2} = \frac{\pi^2}{6}$ give

$$\sum A_n^2 M_n \leqslant \|A\|_2^2 \sup M_n \leqslant \|A\|_2^3\Big(\sum_{m\neq n}\frac{1}{(n-m)^2}\Big)^{\frac{1}{2}} = \frac{\pi}{\sqrt{3}}\|A\|_2^3.$$

We show (4.9). Using (4.7), (2.17),(2.14), we obtain

$$|\nabla_m V_n(h)| \leqslant \frac{8\nu_m h_n^2}{3\pi}\int_{g_n}\frac{v(z)dz}{(n-m)^2} = \frac{4\nu_m}{3}\frac{h_n^2 A_n}{(n-m)^2}, \quad if \quad m \neq n,$$

and using (3.14), we get

$$|\nabla_n V_n(h)| \leqslant \frac{8|h_n|^3}{\pi}\int_{g_n}\frac{v'(z)dz}{z_n - z} \leqslant 8|h_n|^3(1+S_n) \leqslant 9|h_n|^3,$$

which yields (4.9).

Using (3.7),(1.2) we obtain

$$U = \frac{8}{3\pi}\int_g v^3(z)dz \leqslant \frac{8}{3\pi}\|h\|_\infty^2\int_g v(z)dz = \frac{4}{3}\|h\|_\infty^2\|A\|_1 \leqslant \frac{4}{3}\|A\|_1^2,$$

which yields (1.5). ■

Lemma 4.2. *i) Let* $q \in L^2(\mathbb{T})$. *Then*

$$\frac{A_n|h_n|^2}{3} \leqslant \frac{|g_n||h_n|^3}{3\pi} \leqslant V_n \leqslant \frac{4}{3}A_n|h_n|^2, \tag{4.14}$$

$$\frac{1}{3}\sum A_n|h_n|^2 \leqslant V \leqslant \frac{4}{3}\sum A_n|h_n|^2, \tag{4.15}$$

$$\cosh h_n - 1 \leqslant C_0\frac{|g_n|^2}{8}, \qquad C_0 = \cosh\|h\|_\infty, \tag{4.16}$$

$$|h_n| \leqslant \frac{\sqrt{C_0}}{2}|g_n|. \tag{4.17}$$

ii) The estimate (1.6) holds true. Moreover, if $|h_n| = \|h\|_\infty$ *for some* $n \in \mathbb{Z}$, *then*

$$if \qquad C_0 \geqslant 2, \qquad \Rightarrow \quad |g_n| \geqslant 2, \quad |h_n| = \|h\|_\infty \leqslant \frac{\pi}{2}A_n, \tag{4.18}$$

$$C_0 \leqslant C_1 = \max\{2, \cosh\frac{\pi}{2}\|A\|_\infty\}. \tag{4.19}$$

Proof. i) Let $g_n \neq \emptyset$. Assume that $\alpha_n = z_n - z_n^- \geqslant |g_n|/2$, the proof of the case $z_n^+ - z_n \geqslant |g_n|/2$ is similar. Define the function $f_n(t) = t\frac{|h_n|}{\alpha_n}, t = z - z_n^- \in (0,\alpha_n)$. The function $v(z+i0), z \in g_n$ is convex and then $v(z_n^- + t + i0) \geqslant f_n(t), t \in (0,\alpha_n)$, which yields

$$V_n = \frac{8}{3\pi}\int_{g_n} v^3(z,h)dz \geqslant \frac{8}{3\pi}\int_0^{\alpha_n} f_n^3(t)dt = \frac{8}{3\pi}\frac{|h_n|^3}{\alpha_n^3}\int_0^{\alpha_n} t^3dt = \frac{2}{3\pi}|h_n|^3\alpha_n \geqslant \frac{1}{3\pi}|h_n|^3|g_n|.$$

Using $v(z+i0) \leqslant |h_n|$ for all $z \in g_n$, we get $V_n \leqslant \frac{8|h_n|^2}{3\pi}\int_{g_n} v(z,h)dz = \frac{4|h_n|^2}{3}A_n$.

The Taylor formula implies $\Delta(z_n^-) - 1 = \frac{1}{2}\Delta''(\widetilde{z}_n^-)(z_n^- - z_n)^2$ for some $\widetilde{z}_n^- \in (z_n^-, z_n)$.

Using the Bernstein estimates for the bounded exponential type functions we obtain $\sup_{z\in\mathbb{R}}|\Delta''(z)|=\sup_{z\in\mathbb{R}}|\Delta(z)|=C_0$. Then combining the Taylor formula plus the Bernstein estimates we get (4.16), which gives (4.17).

ii) The estimate (4.16) yields $C_0-1\leqslant C_0\frac{|g_n|^2}{8}, \qquad C_0=\cosh\|h\|_\infty$. If $C_0\geqslant 2$, then we deduce that $\frac{C_0}{2}\leqslant C_0\frac{|g_n|^2}{8}$ and thus $|g_n|\geqslant 2$. Thus, the estimate (2.14) gives $|h_n|\leqslant\frac{|h_n||g_n|}{2}\leqslant\frac{\pi}{2}A_n$ and we obtain (4.19).

Using (4.15), (4.17), (4.19), (2.14), we obtain

$$V\leqslant\frac{4}{3}\sum A_n|h_n|^2\leqslant\frac{2}{3}\sqrt{C_0}\sum A_n|h_n||g_n|\leqslant\frac{2\pi}{3}\sqrt{C_1}\|A\|_2^2.$$

Moreover, using (4.15), (2.14), we obtain

$$V\geqslant\frac{1}{3}\sum A_n|h_n|^2\geqslant\frac{\pi}{6}\sum A_n^2=\frac{\pi}{6}\|A\|_2^2,$$

which yields (1.6). ■

The function $V(h)$ is even with respect to each variable $h_n, n\in\mathbb{Z}$, and then $V(h)$ is the function of $h_n^2, n\in\mathbb{Z}$.

Lemma 4.3. *i) Let $h\in\ell^2$ be such that $\|q\|<\frac{1}{8}$. Then for sufficiently small $\varepsilon=(\varepsilon_n)_{n\in\mathbb{Z}}\in\ell^2$ the following estimate holds true:*

$$|V(h+\varepsilon)-V(h)|\leqslant 9\|\varepsilon\|_2. \tag{4.20}$$

ii) The function $V:\ell^2\to[0,\infty)$ has the derivative $\nabla_n V(h)$ for each $n\in\mathbb{Z}$, which is continuous on ℓ^2 and is given by

$$\nabla_m V(h)=\nabla_m V_m(h)-f_m(h), \qquad f_m=\frac{8\nu_m}{3\pi}\int_{g\setminus g_m}\frac{v^3(z)dz}{(z_m-z)^2}, \tag{4.21}$$

$$|f_m|\leqslant 3|\nu_m|\|A\|_\infty S_m. \tag{4.22}$$

Proof. i) Using Lemma 4.1 we get

$$V(h+\varepsilon)-V(h)=\sum_n(V_n(h+\varepsilon)-V_n(h))=\sum_{n,m\in\mathbb{Z}}\int_0^1\nabla_m V_n(h+t\varepsilon)\varepsilon_m dt,$$

where all series converges absolutely. If $\widetilde{h}=h+t\varepsilon$, then estimates (4.9), (2.14) imply

$$|\nabla_n V_n(\widetilde{h})|\leqslant 9|\widetilde{h}_n|^3, \quad and \quad |\nabla_m V_n(\widetilde{h})|\leqslant\frac{4|\nu_m(\widetilde{h})|}{3}\frac{\widetilde{h}_n^2A_n(\widetilde{h})}{(n-m)^2}\leqslant\frac{8}{3\pi}\frac{|\widetilde{h}_m\widetilde{h}_n^4|}{(n-m)^2}, \quad m\neq n,$$

and thus using $\sum_{n>0}\frac{1}{n^2}=\frac{\pi^2}{6}$, we get

$$|V(h+\varepsilon)-V(h)|\leqslant\int_0^1\biggl(\sum_{n\neq m\in\mathbb{Z}}\frac{|\widetilde{h}_m|\widetilde{h}_n^4}{(n-m)^2}|\varepsilon_n|+9\sum_{n\in\mathbb{Z}}|\widetilde{h}_n|^3|\varepsilon_n|\biggr)dt$$

$$\leqslant\|\varepsilon\|_2\int_0^1\biggl(\frac{\pi^2}{3}\|\widetilde{h}\|_2^5+9\|\widetilde{h}\|_2^3\biggr)dt\leqslant 9\|\varepsilon\|_2,$$

since $\|\widetilde{h}\|\leqslant 1$.

ii) Let $e_1=(\delta_{1,n})_{n\in\mathbb{Z}}$ a unit vector in $\ell^p, p\geqslant 1$. Using again Lemma 4.1 we obtain

$$V(h+te_1)-V(h)=\sum_n (V_n(h+te_1)-V_n(h))=\sum_n\int_0^t \nabla_1 V_n(h+\tau e_1)d\tau$$

$$=t\sum_n \nabla_1 V_n(h)+I_0,\qquad I_0=\sum_n\int_0^t \nabla_1\Big(V_n(h+\tau e_1)-V_n(h)\Big)d\tau,$$

where all series converges absolutely. Assume that $I_0=o(t)$ as $t\to 0$. Then Lemma 4.1 gives (4.21).

We show that $I_0=o(t)$ as $t\to 0$. We have

$$I_0=\int_0^t G(t)d\tau,\qquad G(t)=\sum_n b_n(t),\qquad b_n(t)=\nabla_1(V_n(h+\tau e_1)-V_n(h))$$

$$G=G_1+G_2,\qquad G_2=\sum_{-N}^{N} b_n.$$

Estimate (4.9) implies

$$|G_1(t)|\leqslant\sum_{|n|>N}|b_n(t)|\leqslant\sum_{|n|>N}\frac{C}{(1+n^2)}\leqslant\frac{C}{N}$$

for some absolute constant C and $N>1$ large enough. Each function $b_n(t,h)$ is real analytic in $h\in\ell^2$, then $|G_2|=o(t)$ as $t\to 0$.

Similar arguments give that $\nabla_1 V(h)$ is continuous on ℓ^2.

We show (4.22). Using (2.18) and (4.23) we obtain

$$f_{m2}=f_m=\frac{8\nu_m}{3\pi}\int_{g\setminus g_m}\frac{v^3(z)dz}{(z_m-z)^2}\leqslant\frac{8\nu_m\|h\|_\infty^2}{3\pi}\int_{g\setminus g_m}\frac{v(z)dz}{(z_m-z)^2}\leqslant\frac{8\nu_m\|h\|_\infty^2}{3}S_m,$$

and (2.21), (2.19), (2.20), (2.22) imply

$$h_n^2\leqslant(|g_n|^2/4)(1+S_n)^2\leqslant A_n(1+(128)^{-1})^2,\qquad \text{all } n\in\mathbb{Z},$$

which yields (4.22). ■

The function $\widetilde{\omega}_n=\frac{\nabla_n V}{2h_n}$ is well defined and is continuous in ℓ^2, since by Theorem 3.1, the function $\alpha_n=\frac{\nu_n(h)}{2h_n}$ is real analytic.

Lemma 4.4. *i) Let $\alpha_n=\frac{\nu_n(h)}{2h_n}$. Then each component $\widetilde{\omega}_m=\frac{\partial V(h)}{\partial h_m^2}, m\in\mathbb{Z}$ is continuous on ℓ^2 and satisfies for all $h\in\ell^2$:*

$$\widetilde{\omega}_m=\widetilde{\omega}_{m1}-\widetilde{\omega}_{m2},\qquad \widetilde{\omega}_{m1}=\frac{4\alpha_m}{\pi}\int_{g_m}v^2(z)\frac{v'(z)dz}{z_m-z}\geqslant 0,\quad \widetilde{\omega}_{m2}=\frac{4\alpha_m}{3\pi}\int_{g\setminus g_m}\frac{v^3(z)dz}{(z_m-z)^2}\geqslant 0. \tag{4.23}$$

iii) If $\|q\|\leqslant\frac18$, then the components $\widetilde{\omega}_m, m\in\mathbb{Z}$ satisfy

$$\widetilde{\omega}_{m2}\leqslant\frac92|h_n|^2,\qquad \widetilde{\omega}_{m2}\leqslant\frac{3\|A\|_\infty}{2}S_m, \tag{4.24}$$

$$|\widetilde{\omega}_{m1}-2A_m|\leqslant 29A_mS_m\leqslant\frac{A_m}{4}, \tag{4.25}$$

$$|\widetilde{\omega}_m| \leqslant 3A_m + 2\|A\|_\infty S_m, \qquad \|\widetilde{\omega}\|_\infty \leqslant 4\|A\|_\infty, \tag{4.26}$$

$$|\widetilde{\omega}_m - 2A_m| \leqslant 31\|A\|_\infty S_m. \tag{4.27}$$

Proof. i) (4.7) and (4.8) imply the simple estimates (4.23). By Lemma 4.1 and Theorem 3.1, each $\frac{\partial V_n(h)}{\partial h_m^2}, n \in \mathbb{Z}$ is real analytic on ℓ^2 and estimates (4.9) give that each component $\widetilde{\omega}_m = \frac{\partial V(h)}{\partial h_m^2}, m \in \mathbb{Z}$ is continuous on ℓ^2.

ii) Using (4.22), (4.9), (3.7) we obtain (4.24).

Substituting the identity (3.14) into the definition of $\widetilde{\omega}_{m1}$ (see (4.23)) we derive

$$\widetilde{\omega}_{m1} = 4\alpha_m r^2 + I_0 - I, \quad I_0 = \frac{4\alpha_m r^2}{\pi}\int_{g\setminus g_m} \frac{v(t,h)dt}{(t-z_m)^2}, \quad I = \frac{4\alpha_m}{\pi}\int_{g_m} (r^2 - v^2(t))\frac{v'(t)dt}{(z_m - t)}. \tag{4.28}$$

where $r = \frac{|g_m|}{2}$. Estimates (2.18), (2.22) give

$$I_0 = \frac{4\alpha_m r^2}{\pi}\int_{g\setminus g_m} \frac{v(t,h)dt}{(t-z_m)^2} \leqslant 4\alpha_m r^2 S_m \leqslant 4A_m S_m. \tag{4.29}$$

We rewrite the integral I in the form

$$I = \frac{4\alpha_m}{\pi}\int_{g_m} (r^2 - v^2(t))\frac{v'(t)dt}{z_m - t} = I_1 + I_2 + I_3, \quad I_j = \frac{4\alpha_m}{\pi}\int_{g_m} f_j(t)\frac{v'(t)dt}{z_m - t}, \tag{4.30}$$

where

$$r^2 - v^2 = f_1 + f_2 + f_3, \quad f_1 = r^2 - \widetilde{v}, \quad f_2 = \widetilde{v} - v_m^2, \quad f_3 = v_m^2 - v^2, \quad \widetilde{v} = r^2 - (t - z_m)^2$$

and recall that $v_m^2 = r^2 - (t - z_m^0)^2, z_m^0 = \frac{z_m^- + z_m^+}{2}, t \in g_m$. We estimate all $I_j, j = 1,2,3$.

Consider I_1. We have $f_1 = (t - z_m)^2$ and then the integration by parts yields

$$I_1 = \frac{4\alpha_m}{\pi}\int_{g_m} v'(t)(z_m - t)dt = \frac{4\alpha_m}{\pi}\int_{g_m} v(t)dt = 2\alpha_m A_m. \tag{4.31}$$

Consider I_2. Using $f_2 = (z_m - z_m^0)(2t - z_m - z_m^0)$ and (2.26), (2.19) we have

$$|f_2| \leqslant 2|g_m||z_m - z_m^0| \leqslant \frac{|g_m|^3}{2}\dot{M}_m \leqslant 2A_m S_m |g_m|. \tag{4.32}$$

Consider I_3. Using (2.15) we get $f_3 = v_m^2 - v^2 = -v_m^2 Y_m(2 + Y_m)$. Thus, (2.22), (2.19) give

$$|f_3| \leqslant \frac{|g_m|^2}{4}M_m(2 + M_m) \leqslant A_m S_m(2 + S_m). \tag{4.33}$$

Combine (4.30)-(4.31) and using (4.32)-(4.33), (3.14) we have

$$I = 2\alpha_m A_m + I_2 + I_3, \tag{4.34}$$

$$|I_2 + I_3| \leqslant A_m S_m C_T \frac{4}{\pi}\int_{g_m} \frac{v'(t)dt}{z_m - t} \leqslant A_m S_m C_T(4 + 4S_m),$$

where $C_T = 2|g_m| + 2 + S_m$, Using (2.17) and (2.20) we obtain $C_T(4 + 4S_m) \leqslant 11$, thus (4.28), (4.29), (4.34) yield

$$\widetilde{\omega}_{m1} = 2\alpha_m(2r^2 - A_m) + I_0 - I_2 - I_3, \quad |I_2 + I_3| \leqslant 11A_m S_m, \quad I_0 \leqslant 4A_m S_m. \tag{4.35}$$

Consider $\alpha_m(2r^2 - A_m)$, which has the form

$$\alpha_m(2r^2 - A_m) = A_m + I_4, \qquad I_4 = (\alpha_m - 1)A_m + 2\alpha_m(r^2 - A_m)$$

The estimates (2.22) and (2.18) give

$$|r^2 - A_m| \leqslant A_m S_m. \tag{4.36}$$

Estimate (3.21) gives

$$|\alpha_m - 1| \leqslant 5S_m.$$

Thus combine last estimates we obtain

$$|I_4| \leqslant 5S_m A_m + 2S_m A_m = 7S_m A_m,$$

which finally together with (2.20) gives

$$\widetilde{\omega}_{m1} - 2A_m = 2I_4 + I_0 - I_2 - I_3, \qquad |2I_4 + I_0 - I_2 - I_3| \leqslant 29S_m A_m \leqslant A_m/4.$$

This yields (4.25). Combine (4.24), (4.25) and we get (4.27), and additionally using (2.20) we have (4.26). ■

Proof of Theorem 1.1. The estimates (1.5), (1.9) have been proved in Lemma 4.1. Due to [K4] the actions A_n have asymptotics $A_n = |q_n|^2(1 + o(1))$ as $n \to \pm\infty$. Then if $q \in L^{\frac{4}{3}}(\mathbb{T})$, then we deduce that $A \in \ell^2$ and the functional $U(A) < \infty$. ■

Proof of Theorem 1.2. In order to show (1.10) we use the following identity

$$\omega_n = \frac{\partial V}{\partial A_n} = \sum_{m\in\mathbb{Z}} \frac{\partial V}{\partial h_m^2}\frac{\partial h_m^2}{\partial A_n} = \sum_{n\in\mathbb{Z}} X_{n,m}\widetilde{\omega}_m, \quad \widetilde{\omega}_m = \frac{\partial V}{\partial h_m^2}, \quad X_{n,m} = \frac{\partial h_m^2}{\partial A_n}. \tag{4.37}$$

We rewrite the last identity in the short form:

$$\omega = X\widetilde{\omega}, \quad \text{where} \qquad \widetilde{\omega} = (\widetilde{\omega}_m)_{m\in\mathbb{Z}}, \qquad \omega = (\omega_m)_{m\in\mathbb{Z}}, \tag{4.38}$$

and X is an operator in ℓ^2 with coefficients $X_{n,m}$. Note that $X = F^{-1}$, where the operator F is defined in (3.16), (3.17).

Consider the operator F. The estimate (3.20) implies $\|F\| \geqslant 1 - \|F - I_{id}\| \geqslant 1 - \frac{\pi}{64} > \frac{15}{16}$, thus $\|F^{-1}\| \leqslant \frac{16}{15}$. Let $F = I_{id} + B$. Then we obtain

$$\omega = F^{-1}\widetilde{\omega} = \widetilde{\omega} - F^{-1}B\widetilde{\omega} = 2A + f, \qquad \text{where} \quad f = (\widetilde{\omega} - 2A) - F^{-1}B\widetilde{\omega}.$$

Using (4.27) we obtain $|\widetilde{\omega}_m - 2A_m| \leqslant 31\|A\|_\infty S_m$ and (2.20) implies

$$\|\widetilde{\omega} - 2A\|_2 \leqslant 31\|A\|_\infty \|S\|_2 \leqslant 31\|A\|_\infty \frac{\pi^2}{6}\|A\|_2 \leqslant 6\pi^2\|A\|_\infty\|A\|_2. \tag{4.39}$$

Lemma 3.3 yields

$$|(B\widetilde{\omega})_m| \leqslant |(F_{m,m} - 1)\widetilde{\omega}_m| + \sum_{n\neq m} |F_{m,n}\omega_n| \leqslant 5S_m|\widetilde{\omega}_m| + \sum_{n\neq m} \frac{A_n\widetilde{\omega}_n}{(m-n)^2} \leqslant 6\|\widetilde{\omega}\|_\infty S_m.$$

Then (2.20),(4.26) give

$$\|B\widetilde{\omega}\|_2 \leqslant 6\|\widetilde{\omega}\|_\infty \|S\|_2 \leqslant 4\pi^2\|A\|_\infty\|A\|_2. \tag{4.40}$$

Combine (4.39),(4.40) and using $\|F^{-1}\| \leqslant \frac{16}{15}$ we get

$$\|f\|_2 \leqslant \pi^2\|A\|_\infty\|A\|_2(4 + 6\|F^{-1}\|) \leqslant 11\pi^2\|A\|_\infty\|A\|_2,$$

which yields (1.10). ■

Lemma 4.5. *Let* $r \in [0, \pi/2]$ *and* $|z - \pi n| \geqslant r$. *Then*

$$2|\sin z| \geqslant e^{|\operatorname{Im} z|}(1 - e^{-2r}).$$

proof Sufficintely $z \in \overline{\mathbb{C}}_+$. Then

$$2|\sin z|e^{|\operatorname{Im} z|} = 2|e^{iz} \sin z| = |1 - e^{i2z}|$$

The max principle for the $\frac{1}{1-e^{i2z}}$ for the domain

$$\{z \in \overline{\mathbb{C}}_+ : |z - \pi n| \geqslant r, n \in \mathbb{Z}\}$$

and periodic property give that we have to check Lemma for $|z| = r, \operatorname{Im} z \geqslant 0$. Let $w = 1 - e^{i2z}$. We have

$$|z| = r = \frac{1}{2}|\log(1-w)| \leqslant \frac{1}{2}\Big(|w| + \frac{|w|^2}{2} + \frac{|w|^3}{3}...\Big) = \frac{1}{2}\log(1 - |w|)$$

Then $|w| \geqslant 1 - e^{-2r}$.

■

Acknowledgments. The various parts of this paper were written at Mathematical Institute of the Tsukuba Univ., Japan and Ecole Polytechnique, France. The author is grateful to the Institutes for the hospitality. I am grateful to Sergei Kuksin for stimulating discussions and useful comments.

References

[AG] Amour, L.; Guillot, J. Isospectral sets for AKNS systems on the unit interval with generalized periodic boundary conditions. Geom. Funct. Anal. 6 (1996), 127.

[BGGK] D. Battig, B. Grebert, J. Guillot, T. Kappeler, Foliation of phase space for the cubic non-linear Schrödinger equation, Compositio Math., 85(1993), 163-199.

[FT] Faddeev, L. D.; Takhtajan, L. A. Hamiltonian methods in the theory of solitons. Translated from the Russian by A. G. Reyman, Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1987.

[FM] Flashka H., McLaughlin D. Canonically conjugate variables for the Korteveg- de Vries equation and the Toda lattice with periodic boundary conditions. Prog. of Theor. Phys. 55(1976), 438-456.

[GH] Gesztesy, F.; Holden, H. Soliton Equations and their algebro-geometric solutions, Vol. 1. Cambridge University Press, 2003.

[Go] Goluzin, G.M. Geometric theory of functions of a complex variable, Transl. Math. Monogr. , 26 , Amer. Math. Soc. (1969) (Translated from Russian).

[GG] Grebert, B., Guillot, J. Gaps of one-dimensional periodic AKNS systems. Forum Math. 5 (1993), no. 5, 459–504.

[GKP] Grebert, B.; Kappeler, T.; Pöschel, P. Normal form theory for the NLS equation, preprint 2009.

[J] Jenkins A. Univalent functions and conformal mapping. Berlin, Göttingen, Heidelberg, Springer, 1958.

[KP] Kappeler, T.; Pöschel, J. Kdv & Kam. Springer, 2003.

[KK1] Kargaev, P.; Korotyaev, E. The inverse problem for the Hill operator, a direct approach. Invent. Math. 129 (1997), no. 3, 567–593.

[KK2] Kargaev, P.; Korotyaev, E. Effective masses and conformal mappings. Comm. Math. Phys. 169 (1995), no. 3, 597–625.

[KK3] Kargaev, P.; Korotyaev, E. Inverse problems generated by conformal mappings on complex plane with parallel cuts, preprint 2000, ftp://ftp-sfb288.math.tu-berlin.de/pub/Preprints/preprint458.ps.gz

[KK4] Kargaev, P.; Korotyaev, E. Identities for the Dirichlet integral of subharmonic functions from the Cartright class. Complex Var. Theory Appl. 50 (2005), no. 1, 35–50.

[K1] Korotyaev, E. Marchenko-Ostrovki mapping for periodic Zakharov-Shabat systems. J. Differential Equations 175 (2001), no. 2, 244–274.

[K2] Korotyaev, E. Inverse problem and estimates for periodic Zakharov-Shabat systems. J. Reine Angew. Math. 583 (2005), 87–115.

[K3] Korotyaev, E. Metric properties of conformal mappings on the complex plane with parallel cuts. Internat. Math. Res. Notices 1996, no. 10, 493–503.

[K4] Korotyaev, E. A priori estimates for the Hill and Dirac operators, Russ. J. Math. Phys.,15(2008), No. 3, pp. 320-331.

[K5] Korotyaev, E. Hamiltonian and action variables for periodic NLS, in preparation.

[Ku] Kuksin, S. B. Analysis of Hamiltonian PDEs. Oxford Lecture Series in Mathematics and its Applications, 19. Oxford University Press, Oxford, 2000.

[Ku1] Kuksin, S. private communication.

[LS] Levitan B., Sargsjan I. Sturm-Liouville and Dirac operators. Translated from the Russian. Mathematics and its Applications (Soviet Series), 59. Kluwer Academic Publishers Group, Dordericht, 1991, 350 pp.

[L] Löwner K. Untersuchungen über schlichte konforme Abbildung des Einheitskreises, J. Math. Ann., 89(1923), 103-121.

[LM] Lesch, M.; Malamud, M. On the deficiency indices and self-adjointness of symmetric Hamiltonian systems, Journal of Differential Equations 189(2003), 556–615.

[MO] Marchenko V.; Ostrovski I. A characterization of the spectrum of the Hill operator. Math. USSR Sb. 26(1975), 493-554.

[MV1] McKean, H., Vaninsky, K.: Action-angle variables for the cubic Schrödinger equation. Comm. Pure Appl. Math. 50 (1997), no. 6, 489–562.

[MV2] McKean, H. P.; Vaninsky, K. L. Cubic Schrödinger: the petit canonical ensemble in action-angle variables. Comm. Pure Appl. Math. 50 (1997), no. 7, 593–622

[Mi1] Misyura T. Properties of the spectra of periodic and antiperiodic boundary value problems generated by Dirac operators. I,II, Theor. Funktsii Funktsional. Anal. i Prilozhen, (Russian), 30 (1978), 90-101.

[Mi2] Misyura T. Properties of the spectra of periodic and antiperiodic boundary value problems generated by Dirac operators. II, Theor. Funktsii Funktsional. Anal. i Prilozhen, (Russian), 31(1979), 102-109.

[V] Vaninsky, K. L. Symplectic structures and volume elements in the function space for the cubic Schrdinger equation. Duke Math. J. 92 (1998), no. 2, 381–402.

[VN] Veselov, A.; Novikov, S. Poisson brackets and complex tori. Proc. Steklov Inst. Math. 165(1985), 53-65.

[ZS] Zakharov, V.E.; Shabat, A.B. A scheme for integrating nonlinear equations of mathematical physics by the method of the inverse scattering problem I. Funct. Anal. Appl. 8(1974), 226–235.

School of Math., Cardiff University. Senghennydd Road, CF24 4AG Cardiff, Wales, UK. email KorotyaevE@cf.ac.uk, Partially supported by EPSRC grant EP/D054621.